

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8021: Recombinant Human Izumo Sperm-Egg Fusion Protein 4/ IZUMO4 (C-Fc)(Discontinued)

Gene ID: IZUMO4
Gene ID: 113177
Uniprot ID: Q1ZYL8

Description

Source: Human Cells.

MW:50.1kD.

Recombinant Human IZUMO4 is produced by our Mammalian expression system and the target gene encoding His16-His214 is expressed with a Fc tag at the C-terminus. Izumo sperm-egg fusion protein 4 is a sperm membrane protein which plays a key role in the fusion in the mouse. IZUMO4 has an N-terminal domain with significant homology to the N-terminal domain of Izumo.It belongs to the Izumo family . Izumo 4 is a soluble protein expressed in the testis and in other tissues. Izumo domain possesses the ability to form dimers, whereas the transmembrane domain or the cytoplasmic domain or both of Izumo 1 are required for the formation of multimers of higher order.

Product Info

Amount : Fc) / 50 μg

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20 $^{\circ}\text{C},$ though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: HGCLHCHSNFSKKFSFYRHHVNFKSWWVGDIPVSGALLTDWSDDTMKELHLAIPAKITREKLDQVATAVYQM

MDQLYQGKMYFPGYFPNELRNIFREQVHLIQNAIIESRIDCQHRCGIFQYETISCNNCTDSHVACFGYNCESSAQ WKSAVQGLLNYINNWHKQDTSMSLVSPALRCLEPPHLANLTLEDAAECLKQHVDDIEGRMDEPKSCDKTHTC PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.