

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8052: Recombinant Human GADD45gamma/GADD45G (N-6His)(Discontinued)

**Gene ID :** GADD45G **Gene ID :** 10912 **Uniprot ID :** 095257

## **Description**

Source: E.coli. MW:19.28kD.

Recombinant Human GADD45G is produced by our E.coli expression system and the target gene encoding Met1-Glu159 is expressed with a 6His tag at the N-terminus. Growth Arrest and DNA Damage-Inducible Protein GADD45 g (GADD45G) is a nuclear protein which belongs to the GADD45 family. GADD45G is highly expressed in placenta. GADD45G interacts with various proteins whose transcript levels are increased following stressful growth arrest conditions and treatment with DNA-damaging agents. GADD45G responds to environmental stresses by mediating activation of the p38/JNK pathway via MTK1/MEKK4 kinase. GADD45G is also involved in the regulation of growth and apoptosis. GADD45G inhibits cell growth and differentiation by androgens. The mRNA expression is down-regulated in hepatocellular carcinoma.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 150mM NaCl, pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMTLEEVRGQDTVPESTARMQGAGKALHELLLSAQRQGCLTAGVYESAKVLN

VDPDNVTFCVLAAGEEDEGDIALQIHFTLIQAFCCENDIDIVRVGDVQRLAAIVGAGEEAGAPGDLHCILISNPNE

DAWKDPALEKLSLFCEESRSVNDWVPSITLPE

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.