

## 32-8074: Recombinant Human NIP7/KD93 (N-6His)

**Gene :** NIP7  
**Gene ID :** 51388  
**Uniprot ID :** Q9Y221

### Description

Source: E.coli.  
MW :22.6kD.

Recombinant Human NIP7 is produced by our E.coli expression system and the target gene encoding Met1-Thr180 is expressed with a 6His tag at the N-terminus. 60S Ribosome Subunit Biogenesis Protein NIP7 Homolog (NIP7) belongs to the NIP7 family. NIP7 contains one PUA domain, it is essential for the process of proper 27S pre-rRNA and 60S ribosome subunit assembly. NIP7 is a monomer form and interacts with NOL8 and SBDS, and may bind to RNA. In addition, NIP7 is one of the many trans-acting factors required for eukaryotic ribosome biogenesis, which interacts with nascent pre-ribosomal particles and dissociates as they complete maturation and are exported to the cytoplasm.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 100mM NaCl, pH 8.0 .  
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.  
**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MGSSHHHHHHSSGLVPRGSHMRPLTEEETRVMEFKIYIGENLQLLVDRPDGTYCFRLHNDRVVYVSEKIMK  
LAANISGDKLVSLGTFCFGKFTKTHKFRHLHVTALDYLAPEYAKYKVVWIKPGAEQSFLYGNHVLKSGGLGRITENTSQY  
QGVVVYSMADIPLGFGVAAKSTQDCRKVDPMAIVVFHQADIGEYVRHEETLT

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.