

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8075: Recombinant Human Isopentenyl Pyrophosphate Isomerase 2//IPPI2/IDI2 (N-6His)

Gene ID: 91734
Uniprot ID: Q9BXS1

## **Description**

Source: E.coli. MW :28.9kD.

Recombinant Human IPP Isomerase 2 is produced by our E.coli expression system and the target gene encoding Met1-Val227 is expressed with a 6His tag at the N-terminus. Isopentenyl Pyrophosphate Isomerase 2 (IDI2) belongs to the IPP isomerase type 1 family. Both isozymes, IDI1 and IDI2 are localized to the peroxisome by a PTS1-dependent pathway. IDI2 is expressed in skeletal muscle, which contains one nudix hydrolase domain. IDI2 binds one magnesium per subunit. IDI2 catalyzes the 1,3-allylic rearrangement of the homoallylic substrate isopentenyl (IPP) to its highly electrophilic allylic isomer, dimethylallyl diphosphate (DMAPP). It is reported that IDI2 is regulated independently from IDI1, by a mechanism that may involve PPAR-a.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Supplied as a 0.2 μm filtered solution of 20mM TrisHCl, 1mM DTT, 0.1mM PMSF, pH 8.0. Storage condition: Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMSDINLDWVDRRQLQRLEEMLIVVDENDKVIGADTKRNCHLNENIEKGLLHR

AFSVVLFNTKNRILIQQRSDTKVTFPGYFTDSCSSHPLYNPAELEEKDAIGVRRAAQRRLQAELGIPGEQISPEDI VFMTIYHHKAKSDRIWGEHEICYLLLVRKNVTLNPDPSETKSILYLSQEELWELLEREARGEVKVTPWLRTIAERF

LYRWWPHLDDVTPFVELHKIHRV

## **Application Note**

**Endotoxin :** Less than  $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.