

32-8099: Recombinant Human Nuclear Transcription Factor Y Subunit α /NFYA (N-GST)

Gene : NFYA
Gene ID : 4800
Uniprot ID : P23511

Description

Source: E.coli.
MW :60.58kD.

Recombinant Human Nuclear TF Y subunit α is produced by our E.coli expression system and the target gene encoding Met1-Ser318 is expressed with a GST tag at the N-terminus. Nuclear Transcription Factor Y Subunit α (NFYA) is a member of the NFYA/HAP2 subunit family. NFYA functions as a heterotrimeric transcription factor, which is composed of three components, NF-YA, NF-YB and NF-YC, binds to CCAAT motifs in the promoter regions in a variety of genes. NFYA forms a highly conserved transcription factor which stimulates the transcription of various genes by recognizing and binding to a CCAAT motif in promoters, for example in type 1 collagen, albumin and beta-actin genes.

Product Info

Amount : 10 μ g / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MSPILGYWKIKGLVQPTRLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPYYIDGDVKLTQSMAIRYIA
DKHNMLGGCPKERAIEISMLEGAVLDIRYGVSRIAYSKDFETLKVDFLSKLEMLKMFEDRLCHKTYLNGDHVTH
PDFMLYDALDVVLYMDPMCLDAFPKLVCFKKRIEAIPIQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPR
GSPEFMEQYTANSNSSTEQIVVQAGQIQQQVQGQPLMVQVSGGQLITSTGQPIMVQAVPGGQQTIMQVPVS
GTQGLQQIQLVPPGQIQIQGGQAVQVQGGQQTQIIIQQPQTAVTAGQTQTQQQIAVQGGQVAQTAEGQTI
VYQPVNADGTILQQVTVPVSGMITIPAASLAGAQIVQTGANTNTTSSGQGTVTLPVAGNVVNSGGMVMMP
GAGSVPAIQRIPLPGAEMLEEEPLYVNAKQYNRILKRRQARAKLEAEGKIPKERRKYLHESRHRHAMARKRREG
GRFFSPKEKDSPHMQDPNQADEEAMTQIRVS

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.