

32-8129: Recombinant Mouse C-X-C Motif Chemokine 2/CXCL2/MIP-2

Gene : Cxcl2
Gene ID : 20310
Uniprot ID : P10889

Description

Source: E.coli.
MW :7.9kD.

Recombinant Mouse C-X-C motif chemokine 2 is produced by our E.coli expression system and the target gene encoding Ala28-Asn100 is expressed. C-X-C motif chemokine 2 (CXCL2,MIP-2) belongs to the intercrine alpha (chemokine CxC) family. It was originally identified as a heparin-binding protein secreted from a murine macrophage cell line in response to endotoxin stimulation. The expression of mouse MIP-2 is stimulated by endotoxin. The mouse MIP-2 shares approximately 63% aa sequence identity with murine KC, another mouse alpha chemokine, which is induced by PDGF. It has been suggested that mouse KC and MIP-2 are the homologs of the human GROs and rat CINC_s. Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst. The expression of MIP-2 was found to be associated with neutrophil influx in pulmonary inflammation and glomerulonephritis, suggesting that MIP-2 may contribute to the pathogenesis of inflammatory diseases.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl,pH8.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AVVASELRCQCLKTLPRVDFKNIQSLSVTPPGPHCAQTEVIATLKGQKVCCLDPEAPLVQKIIQKILNKGKAN

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.