

## 32-8149: Recombinant Human Allograft Inflammatory Factor 1/AIF1 (C-6His)(Discontinued)

**Gene :** AIF1  
**Gene ID :** 199  
**Uniprot ID :** P55008

### Description

Source: E. coli.  
MW :17.7kD.

Recombinant Human Allograft Inflammatory Factor 1 is produced by our E.coli expression system and the target gene encoding Ser2-Pro147 is expressed with a 6His tag at the C-terminus. Allograft Inflammatory Factor 1 (AIF1) contains two EF-hand domains and exists as a homodimer. AIF1 can be detected in T-lymphocytes and peripheral blood mononuclear cells. AIF1 functions as actin-binding protein that enhances membrane ruffling and RAC activation and can enhance the actin-bundling activity of LCP1. In addition, AIF1 plays a role in RAC signaling and in phagocytosis and may also in macrophage activation and function. AIF1 promotes the proliferation of vascular smooth muscle cells and of T-lymphocytes and plays a role in vascular inflammation.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** SQTRDLQGGKAFGLLKAQQEERLDEINKQFLDDPKYSSDEDLPSKLEGFKEKYMFDLNGNGDIDIMSLKRMLEKLGVPKTHLELKKLIGEVSSGSETFSYPDFLRMMLGKRSAILKMILMYEEKAREKEKPTGPPAKKAISELPLEHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.