

32-8168: Recombinant Human SMAD Family Member 4/SMAD4/DPC4 (C-6His)(Discontinued)

Gene : SMAD4
Gene ID : 4089
Uniprot ID : Q13485

Description

Source: E. coli.
MW :61.5kD.

Recombinant Human SMAD4 is produced by our E.coli expression system and the target gene encoding Met1-Asp552 is expressed with a 6His tag at the C-terminus. SMAD Family Member 4 (SMAD4) is a cytoplasmic protein that belongs to the Dwarfing/SMAD family. SMAD4 contains one MH1 (MAD homology 1) domain and one MH2 (MAD homology 2) domain. It is the component of the heterotrimeric SMAD2/SMAD3-SMAD4 complex that forms in the nucleus and is required for the TGF-mediated signaling. SMAD4 promotes binding of the SMAD2/SMAD4/FAST-1 complex to DNA and provides an activation function required for SMAD1 or SMAD2 to stimulate transcription. SMAD4 may act as a tumor suppressor. It positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator. Mutations or deletions in SMAD4 have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl,pH8.0.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : MDNMSITNTPTSNDACLIVHSLMCHRQGGSETFKRAIESLVKCLKKDELDSLITAITTNGAHPKCVTIQR
TLDGRLQVAGRKGFPHVIYARLWRWPDHLKNEKLVKCYQYAFDLKCDSCVNPYHYERVVSPGIDLSGLTLQ
SNAPSSMMVKDEYVHDFEGQPSLSTEGHSIQTIQHPPSNRASTETYSTPALLAPSES NATSTANFPNIPVASTSQ
PASILGGSHSEGLLQIASGPQPGQQNGFTGQPATYHHNSTTTWTGSRTPYTPNLPHHQNGHLQHHPMP
HPGHYWPVHNELAFQPPISNHPAPEYWCSIAYFEMDVQVGETFKVPSSCPVTVDGYVDPSSGDRFCLGQLSN
VHRTEAIERARLHIGKGVQLECKGEGDVWVRLSDHAVFVQSYLDREAGRPGDAVHKIYPSAYIKVFDLRQ
CHRQMQQQAATAQAAAAAQA AVAGNIPGPGSVGGIAPAI SL SAAAGIVDDLRLRLCILRMSFVKGWGPDYP
RQSIKETPCWIEIHLRALQLLDEVLHTMPIADPQPLDLEHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.