

## 32-8189: Recombinant Human Zinc Finger Protein 762/ZFN762/ZIK1 (N-T7 tag)(Discontinued)

**Gene :** ZIK1  
**Gene ID :** 284307  
**Uniprot ID :** Q3SY52

### Description

Source: E. coli.  
MW :44.6kD.

Recombinant Human Zinc Finger Protein 762 is produced by our E.coli expression system and the target gene encoding Met1-Cys384 is expressed with a T7 tag at the N-terminus. Zinc Finger Protein Interacting with Ribonucleoprotein K (ZIK1) is a 487 amino acid nuclear protein that belongs to the Krueppel C2H2-Type Zinc-Finger Protein family. ZIK1 has nine C2H2-type zinc fingers and a KRAB domain. This protein is expressed at high levels in the gastric glands and at low levels in the colon and small intestine. It has been shown that ZIK1 is a transcriptional repressor that interacts with the Heterogeneous Nuclear Ribonucleoprotein Particle K Protein (HNRPK).

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM Tris, pH 7.5.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MASMTGGQQMGRGSM AAAALRAP TQVTVSPETHMDLTGKCVTFEDIAIYFSQDEWGLLDEAQRLLYLEVMLE  
NFALVASLGC GHGTEDEETPSDQNVSVGVSQSKAGSSTQKTQSC EMCVPVLKDILHLADLP GQKPYLVGECTN  
HHQH QKHHS AKKSLKRDMDRASYVKCC LFCMSLKPFRKWEVGKDL PAMLRLLRSLVFP GGKPGTIT ECGEDI  
RSQKSHYKSGECGKASRHKHTPVYHPRVYTGKLYECSKCGKAFRGKYSLVQHQRVHTGERPWECNECGKFF  
SQTSHLNDHRR IHTGERPYECSECGKLF RQNSSLV DHQKIHTGARPYECSQCGKSFSQKATLVKHQRVHTGER  
PYKCGECGNSFSQSAILNQHRR IHTGAKPYECGQC

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.