

32-8213: Recombinant Human MHC Class I Polypeptide-Related Sequence A/MICA (C-Fc)

Gene : MICA
Gene ID : 100507436
Uniprot ID : Q29983

Description

Source: Human Cells.
MW :59.9kD.

Recombinant Human MHC Class I-related Protein A is produced by our Mammalian expression system and the target gene encoding Ala23-Glu308 is expressed with a Fc tag at the C-terminus. MHC class I polypeptide-related sequence A, also known as MIC-A, PERB11.1 and MICA, is a single-pass type I membrane protein which belongs to the MHC class I family of MIC subfamily. MICA contains one Ig-like C1-type domain and is expressed on the cell surface, although unlike canonical class I molecules does not seem to associate with beta-2-microglobulin. It is thought that MICA functions as a stress-induced antigen that is broadly recognized by NK cells, NKT cells, and most of the subtypes of T cells. MICA is the ligand for NK cell activating receptor KLRK1/NKG2D. MICA seems to have no role in antigen presentation. MICA leads to cell lysis by binding to KLRK1.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AEPHSLRYNLTVLSWDGVSQSGFLTEVHLDGQPFLRCRDRQKCRAPQGQWAEDVLGNKTWDRETRDLTGN GKDLRMTLAHIKDQKEGLHSLQEIRVCEIHEDNSTRSSQHFYYDGELFSLQNETEETWMPQSSRAQTLAMNV RNFLKEDAMKTKTHYHAMHADCLQELRRYLKSGVLRRTVPPMVNTRSEASEGNITVTCRASGFYPWNITLS WRQDGVSLSHDTQQWGDVLPDNGTYQTWVATRICQGEQRFTCYMEHSGNHSTHPVPSGKVLVLQSHW QVDDIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYV DGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPP SREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVDFCS VMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.