

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8433: Recombinant Human IL-17 Receptor A/IL-17RA/CD217 (C-Fc)

Gene ID: 23765 Uniprot ID: Q96F46

Description

Source: Human Cells. MW :60.6kD.

Recombinant Human Interleukin-17 receptor A is produced by our Mammalian expression system and the target gene encoding Leu33-Trp320 is expressed with a Fc tag at the C-terminus. Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure.

Product Info

Amount: $50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LRLLDHRALVCSQPGLNCTVKNSTCLDDSWIHPRNLTPSSPKDLQIQLHFAHTQQGDLFPVAHIEWTLQTDASI

LYLEGAELSVLQLNTNERLCVRFEFLSKLRHHHRRWRFTFSHFVVDPDQEYEVTVHHLPKPIPDGDPNHQSKNF LVPDCEHARMKVTTPCMSSGSLWDPNITVETLEAHQLRVSFTLWNESTHYQILLTSFPHMENHSCFEHMHHIP APRPEEFHQRSNVTLTLRNLKGCCRHQVQIQPFFSSCLNDCLRHSATVSCPEMPDTPEPIPDYMPLWVDDIEGR MDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL

HNHYTQKSLSLSPGK

Application Note

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.




Figure 1: Coomassie staining of IL-17RA under denaturing conditions. 5ug and 10ug were loaded.