

32-8436: Recombinant Human CD40 Ligand/CD40L/TNFSF5

Gene : CD40LG

Gene ID : 959

Uniprot ID : P29965

Description

Source: Human Cells.

MW :16.2kD.

Recombinant Human TNFSF5 is produced by our Mammalian expression system and the target gene encoding Met113-Leu261 is expressed. CD40 Ligand (CD40LG) is a type II transmembrane glycoprotein that belongs to the TNF superfamily. Like other TNF superfamily members, CD40LG exists as a trimer in membrane bound and soluble form, both of which are bioactive. CD40LG is a ligand for CD40; its ligation also initiates signal transduction in CD40LG expressing cells. CD40LG is a differentiation antigen that is expressed on the surface of T-cells. It stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines. CD40LG has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also co-stimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma, TNF-alpha, and IL2.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 200mM NaCl, 0.1mM EDTA, pH 7.0.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : MQKGDQNPQIAAHVISEASSKTTSVLQWAEKGYTMSNNLVLENGKQLTVKRQGLYYIYAQVTFCSNREASS QAPFIASLCLKSPGRFRERILLRAANTHSSAKPCGQQSIHLGGVFELQPGASVFNVTDPQSQVSHGTGFTSFGLLK L

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.