

32-8448: Recombinant Human Hyaluronan-Binding Protein 2/HABP2 (C-6His)

Gene : HABP2
Gene ID : 3026
Uniprot ID : Q14520

Description

Source: Human Cells.
MW :36.2kD.

Recombinant Human Hyaluronan-binding Protein 2 is produced by our Mammalian expression system and the target gene encoding Met1-Gln279 is expressed with a 6His tag at the C-terminus. Hyaluronan-binding protein 2(HABP2) is an extracellular serine protease which binds hyaluronic acid. It secreted as an inactive single-chain precursor and is then activated to a heterodimeric form, which consists of a 50 kDa heavy and a 27 kDa light chain linked by a disulfide bond. HABP2 is involved in cell adhesion, it can cleave the alpha-chain at multiple sites and the beta-chain between 'Lys-53' and 'Lys-54' , but not the gamma-chain of fibrinogen. As a result of this, it does not initiate the formation of the fibrin clot and does not cause the fibrinolysis directly. It does not cleave prothrombin and plasminogen but converts the inactive single chain urinary plasminogen activator to the active two chain form, activates coagulation factor VII.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 .
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MFARMSDLHVLLLMALVGKTACGFLMSLLES LDPDWTPDQYDYSYEDYNQEENTSS TLTHAENPDWYTED QADPCQPNPCEHGGDCLVHGSTFTCSCLAPFSGNKCQKQVQNTCKDNPCGRGQCLITQSPYYRCVCKHPYTG PSCSQVVPVCRPNPCQNGATCSRHKRRSKFTACPDQFKGKFCEIGSDDCYVGDGYSYRGKMNRTVNQHAC LYWNSHLLLQENYNMFMEDAETHGIGEHNFCRNPDADEKPWCFIKVTNDKVKWEYCDVSACSAQAVDHHHH HH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.