

## 36-1516: Monoclonal Antibody to NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker)(Clone : SPM299)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM299
<b>Application :</b>	IHC,FACS,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	NGFR
<b>Gene ID :</b>	4804
<b>Uniprot ID :</b>	P08138
<b>Format :</b>	Purified
<b>Alternative Name :</b>	NGFR,TNFRSF16
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	NGFR from A875 melanoma cells

### Description


It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). Its epitope spans in aa 1-160 of extracellular domain of NGFR/NTR. NGF-receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain. NGF is important for the development, differentiation, and survival of variety of neuronal and non-neuronal cells. Its action is mediated by binding two distinct receptors, the high affinity p140 and low affinity p75.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Melanoma stained with NGFR Monoclonal Antibody (SPM299).