

32-8478: Recombinant Human Leukocyte Elastase Inhibitor/Serpin B1/SERPINB1 (C-6His)

 Gene :
 SERPINB1

 Gene ID :
 1992

 Uniprot ID :
 P30740

Description

Source: Human Cells.

MW :43.8kD.

Recombinant Human Serpin B1/Leukocyte elastase inhibitor is produced by our Mammalian expression system and the target gene encoding Met1-Pro379 is expressed with a 6His tag at the C-terminus. SERPINB1 is a member of the serpin family and Ov-serpin subfamily. As protease inhibitors, serpins have an array of functions including regulating blood coagulation, fibrinolysis, the complement pathway, angiogenesis, inflammation, tumor suppression, extracellular matrix remodeling, and cell motility. SERPINB1 regulates the activity of the neutrophil proteases elastase, cathepsin G, proteinase-3, chymase, chymotrypsin, and kallikrein-3. Reactive bond 1 of SerpinB1 is specific for reaction with chymotrypsin-like protease such as cathepsin G, chymotrypsin or chymase. Reactive bond 2 of SerpinB1 is specific for reaction, sERPINB1 also functions as a potent intracellular inhibitor of granzyme H.

Product Info

Amount :	10 μg / 50 μg
Content :	Lyophilized from a 0.2 μ m filtered solution of 20mM PB,150mM NaCl, pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	MEQLSSANTRFALDLFLALSENNPAGNIFISPFSISSAMAMVFLGTRGNTAAQLSKTFHFNTVEEVHSRFQSLNA DINKRGASYILKLANRLYGEKTYNFLPEFLVSTQKTYGADLASVDFQHASEDARKTINQWVKGQTEGKIPELLAS GMVDNMTKLVLVNAIYFKGNWKDKFMKEATTNAPFRLNKKDRKTVKMMYQKKKFAYGYIEDLKCRVLELPYQG EELSMVILLPDDIEDESTGLKKIEEQLTLEKLHEWTKPENLDFIEVNVSLPRFKLEESYTLNSDLARLGVQDLFNSS KADLSGMSGARDIFISKIVHKSFVEVNEEGTEAAAATAGIATFCMLMPEENFTADHPFLFFIRHNSSGSILFLGRF SSPVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.




Figure 1: SDS-PAGE analysis of purified Recombinant Serpin B1 His Tag. 5 μg protein was run on a 4-20% SDS-PAGE gel followed by Coomassie blue staining

For Research Use Only. Not for use in diagnostic/therapeutics procedures.