

## 32-8502: Recombinant Human Interleukin-12 Subunit beta/IL-12 p40/IL-12B(Discontinued)

**Gene :** IL12B  
**Gene ID :** 3593  
**Uniprot ID :** P29460

### Description

Source: Human Cells.  
MW :34.6kD.

Recombinant Human Interleukin-12 subunit beta is produced by our Mammalian expression system and the target gene encoding Ile23-Ser328 is expressed. Interleukin-12 subunit beta (IL-12B) belongs to the type I cytokine receptor family. It contains 1 fibronectin type-III domain and 1 Ig-like C2-type domain. IL-12B is a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. IL-12 is a disulfide-linked heterodimer composed of the 40 kD cytokine receptor encoded by IL12B and a 35 kD subunit encoded by IL12A. IL12 is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. It has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** IWELKKDVYVVELDWYPDAPGEMVLTCDTPEEDGITWTLQDQSSEVLGSGKTLTIQVKEFGDAGQYTCHKGGELVLSHLLLLLHKKEDGIWSTDILKDQKEPKNKTFLRCEAKNYSGRFTCWWLTTISTDLTFSVKSSRGSSDPQGVTCGAATLSAERVGRDNKEYEYSVEQCEDSACPAEESLPIEVMVDAVHKLKYENYTSFFIRDIKPDPPKNLQLKPLKNSRQVEVSWEYPDTWSTPHSYFSLTFCVQVQGKSKREKDRVFTDKTSATVICRKNASISVRAQDRYSSSWSEWASVPCS

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.