

## 32-8504: Recombinant Human C-X-C Motif Chemokine 4/CXCL4/PF4 (C-6His)

**Gene :** PF4  
**Gene ID :** 5196  
**Uniprot ID :** P02776

### Description

Source: Human Cells.  
MW :8.5kD.

Recombinant Human C-X-C Motif Chemokine 4 is produced by our Mammalian expression system and the target gene encoding Glu32-Ser101 is expressed with a 6His tag at the C-terminus. Human Chemokine (C-X-C motif) Ligand 4 (CXCL4) is expressed in megakaryocytes and stored in the alpha-granules of platelets. CXCL4 contains several heparin-binding sites at the C-terminal region and binds heparin with high affinity. The active CXCL4 protein is a tetramer. Human and mouse CXCL4 share 64% sequence identity. CXCL4 is chemotactic for neutrophils, fibroblasts and monocytes and plays a critical role in inflammation and wound repair. CXCL4 functions via a splice variant of the chemokine receptor CXCR3, known as CXCR3B. The major physiologic role of CXCL4 appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. In contrast to other CXC chemokines, CXCL4 lacks chemotactic activity for polymorphonuclear granulocytes.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** EA EEDGDLQCLCVKTT SQVRPRHITSLEVIKAGPHCPTAQLIATLKNRGIKLDLQAPLYKKI IKLLSHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.