

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8558: Recombinant Mouse NKG2D Ligand 1/NKG2DL/ULBP1 (C-6His)(Discontinued)

Gene ID: 77777
Uniprot ID: Q8HWA3

Description

Source: Human Cells. MW :22.3kD.

Recombinant Mouse NKG2D ligand 1 is produced by our Mammalian expression system and the target gene encoding Pro26-Thr211 is expressed with a 6His tag at the C-terminus. Mouse ULBP1, also known as RAET1I and NKG2DL1, is a member of the ULBP/RAET1 gene family. ULBP1 plays an important role in immune responses, especially in cancer and infectious diseases, and is well-known to bind to NKG2D together with at least ULBP 2 and 3. These proteins are distantly related to major histocompatibility class I (MHC I) molecules, possessing the alpha 1 and alpha 2 Ig-like domains, but lacking the alpha 3 domain. Unlike MHC Class I, they have no capacity to bind peptide or interact with beta2-microglobulin. It can activate multiple signaling pathways in primary NK cells, gamma delta T cells, and CD8+ alpha beta T cells, resulting in the production of cytokines and chemokines.ULBP1 is expressed in wide range of tissues including heart, brain, lung, liver, bone marrow and some tumor cells, T-cells, B-cells, As an unconventional member of the MHC class I family, ULBP1 is able to interact with soluble CMV glycoprotein UL16 in CMV infected cells. The interaction with UL16 blocked the interaction with the NKG2D receptor, and thus might escape the immune surveillance. Furthermore, UL16 also causes ULBP1 to be retained in the ER and cis-Golgi apparatus so that it does not reach the cell surface. The ULBP1 regulation may have implications for development of new therapeutic strategies against cancer cells.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: PRIEETASLCNIYKVNRSESGQHSHEVQGLLNRQPLFVYKDKKCHAIGAHRNSMNATKICEKEVDTLKDGIDIFK

GLLLHIVQETNTTGKPLTLQAEVCGQYEVDKHFTGYAIVSLNGKNIFRVDTSTGNWTQLDHEFEKFIEMCKEDK

VLAAFLKKTTEGDCRTWLDELMLHWKEHLEPAGSFSTVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.