

32-8562: Recombinant Human Lymphocyte Activation Gene 3/LAG-3/CD223 (C-6His)(Discontinued)

Gene : LAG3
Gene ID : 3902
Uniprot ID : P18627

Description

Source: Human Cells.

MW :45.5kD.

Recombinant Human LAG-3 is produced by our Mammalian expression system and the target gene encoding Leu23-Gly434 is expressed with a 6His tag at the C-terminus. Human Lymphocyte activation gene 3 protein(LAG3) is a member of immunoglobulin (Ig) superfamily. LAG3 contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. LAG3 is involved in lymphocyte activation and can bind to HLA class-II antigens. It is selectively expressed in activated T and NK cells. LAG3 has a negative regulatory function in T cells and acts as as a new marker of T cell induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signaling. It can lead to increased antigen-specific T-cell responses in vivo. LAG-3 has higher affinity to MHC class II than CD4.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : LQPGAIEVPPVWAQEGAPALPCSPTIPLQDLSLLRRAGVTWQHQPDSGPPAAAPGHPLAPGHPAAPPSSWGP
RPRRYTVLSVPGGLRSGRLPLQPRVQLDERGRQRGDFSLWLRPARRADAGEYRAAVHLRDRALSCRLRLRL
GQASMTASPPGSLRASDWVILNCSFSRPDRPASVHWFNRNGQGRVPVRESPPHHLAESFLFLPQVSPMDSGP
WGCILTYRDGFNVSIMYNLTVLGLPPTPLTVYAGAGSRVGLPCRLPAGVGTRESFLTAKWTPPGGGPDLLVTGD
NGDFTLRLEDVVSQAQAGTYTCHIHLEQQQLNATVTLAIITVTPKSFSGPSLGLKLLCEVTPVSGQERFVWSSLDL
PSQRSFSGPWLEAQEAQLLSQPWQCQLYQGERLLGAAVYFTELSSPGHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.