

32-8570: Recombinant Mouse PD-L2/B7-DC/CD273 (C-Fc)

Gene : Pcd1lg2

Gene ID : 58205

Uniprot ID : Q9WUL5

Description

Source: Human Cells.

MW :49.7kD.

Recombinant Mouse Programmed Cell Death 1 Ligand 2 is produced by our Mammalian expression system and the target gene encoding Leu20-Arg219 is expressed with a Fc tag at the C-terminus. Programmed cell death 1 ligand 2 (PD-L2), also known as butyrophilin B7-DC or PDCD1 ligand 2, belongs to the member of B7 family which can regulate the activation and tolerance of T cells. PD-L2 is one ligand for Programmed cell death 1(PD-1), and the other is PD-L1. These two ligands shares 34% aa sequence identity. Mouse PD-L2 gene encodes a 273 amino acids (aa) protein with a putative 19 aa signal peptide, a 201 aa extracellular region , a 21 aa transmembrane domain and a 32 aa cytoplasmic region. The mouse PD-L2 gene is highly expressed in heart, placenta, pancreas, lung and liver while expressed weakly in spleen, lymph nodes and thymus. Besides, the expression of PD-L2 gene can be induced on dendritic cells grown from peripheral blood mononuclear cells under CSF2 and IL4/interleukin-4 treatment, and up-regulated by IFNG/IFN-gamma stimulation in monocytes. PD-L2 usually functions in a PDCD1-independent manner and is involved in regulating costimulatory signal which is essential for T-cell proliferation and IFNG production. Recent studies demonstrate that the expression of PD-L2 on the tumor cells promotes CD8 T cell-mediated rejection of tumor cells, at both the induction and effector phase of antitumor immunity. Moreover, PD-L2 binds to PD-1 cells and enhances T cell killing in a PD-1-independent mechanism.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : LFTVTAPKEVYTVDVGSSVSLECDFDRRECTELEGRASLQKVENDTSLQSERATLLEEQLPLGKALFHIPSVQV
RDSGQYRCLVICGAAWDYKYLTVKVKASYMRIDTRILEVPGTGEVQLTCQARGYPLAEVSWQNVSPANTSHI
RTPEGLYQVTSVLRKLPQPSRNFSCMFVNAHMKELTSIIDPLSRMEPKVPRVDDIEGRMDEPKSCDKTHTCPP
CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV
VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI
AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.