

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8581: Recombinant Mouse Interleukin-16/IL-16

**Gene ID:** 16170 **Uniprot ID:** 054824

## **Description**

Source: E.coli. MW :14.5kD.

Recombinant Mouse Interleukin-16 is produced by our E.coli expression system and the target gene encoding Ser1205-Ser1322 is expressed with a 6His tag at the N-terminus. Mouse interleukin-16(IL-16) is a single chain non-glycosylated polypeptide. IL-16 is widely expressed in human tissues including spleen, thymus, lymph nodes, peripheral leukocytes, bone marrow and cerebellum. IL-16 plays an important role instimulating a migratory response in CD4+ lymphocytes, monocytes, and eosinophils,inducing T-lymphocyte expression of interleukin 2 receptor. It was originally identified as a CD8+ T cell-derived chemoattractant for CD4+ cells. In addition to its chemotactic properties, IL-16 has also been shown to suppress HIV-1 replication in vitro and appears to be involved in transcriptional regulation of SKP2 and is probably part of a transcriptional repression complex on the core promoter of the SKP2 gene. It may act as a scaffold for GABPB1 (the DNA-binding subunit the GABP transcription factor complex) and HDAC3 thus maintaining transcriptional repression and blocking cell cycle progression in resting T-cells.

## **Product Info**

**Amount:** 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl,pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMSAASASAASDISVESKEATVCTVTLEKTSAGLGFSLEGGKGSLHGDKPLTIN

RIFKGTEOGEMVOPGDEILOLAGTAVOGLTRFEAWNVIKALPDGPVTIVIRRTSLOCKOTTASADS

## **Application Note**

**Endotoxin :** Less than  $0.1 \text{ ng/}\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.