

## 32-8586: Recombinant Human Thymic Stromal Lymphopoietin/TSLP(Discontinued)

**Gene :** TSLP  
**Gene ID :** 85480  
**Uniprot ID :** Q969D9

### Description

Source: E.coli.  
MW :15.1kD.

Recombinant Human Thymic stromal lymphopoietin is produced by our E.coli expression system and the target gene encoding Tyr29-Gln159 is expressed. Thymic stromal lymphopoietin (TSLP) is a novel member of the hemopoietic cytokine family that promotes the development of B cells and shares overlapping activity with IL-7. The human TSLP protein comprises a 28 amino acids (aa) signal sequence and 131 aa mature region. Human TSLP has two isoforms lTSLP (159 aa) and sTSLP (63 aa) produced by alternative splicing. lTSLP (159aa) is expressed in a number of tissues including heart, liver and prostate, and sTSLP (63aa) is predominantly expressed in keratinocytes of oral mucosa, skin and in salivary glands. In aa sequence level, Human TSLP displays about 43% identity with mouse TSLP. Thymic stromal lymphopoietin (TSLP) is a cytokine that functions mainly on myeloid cells; it induces the release of T cell-attracting chemokines from monocytes and enhances the maturation of CD11c(+) dendritic cells. TSLP has proliferative effects on the myeloid cell line and may initiate asthma or atopic dermatitis responses by directly activating mast cells. TSLP signals cells via the interleukin-7 receptor- $\alpha$  chain (IL-7Ra), shared with IL-7, together with the TSLP receptor (TSLPR) subunit. Recent studies indicate that TSLP and its receptor are novel therapeutic targets for rheumatoid arthritis, for increased intraarticular TSLP concentrations in patients has caused chemotaxis and activation of arthritogenic T cells.

### Product Info

**Amount :** 10  $\mu$ g / 50  $\mu$ g  
**Content :** Lyophilized from a 0.2  $\mu$ m filtered solution of PBS,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** YDFTNCDFEKIKAAYLSTISKDLITYMSGTKSTEFNNTVSCSNRPHCLTEIQSLTFNPTAGCASLAKEMFAMKTKA  
ALAIWCPGYSETQINATQAMKKRRKRKVTNNKCLEQVSQQLQGLWRRFNRPLLKQQ

### Application Note

**Endotoxin :** Less than 0.1 ng/ $\mu$ g (1 IEU/ $\mu$ g) as determined by LAL test.