

## 32-8621: Recombinant Mouse OX40/TNFRSF4/CD134 (N-Fc)

**Gene :** Tnfrsf4

**Gene ID :** 22163

**Uniprot ID :** P47741

### Description

Source: Human Cells.

MW :47.4kD.

Recombinant Mouse OX40 is produced by our Mammalian expression system and the target gene encoding Val20-Pro211 is expressed with a Fc tag at the N-terminus. OX40, also termed CD134 and TNFRSF4, is a T cell co-stimulatory molecule of the TNF receptor superfamily which plays a key role in the survival and homeostasis of effector and memory T cells. OX40 is expressed on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. The interaction between OX40 and OX40 ligand (OX40L) will occur when activated T cells bind to professional antigen-presenting cells (APCs). The T-cell functions, including cytokine production, expansion, and survival, are then enhanced by the OX40 costimulatory signals. OX40 signals are critical for controlling the function and differentiation of Foxp3+ regulatory T cells. OX40-OX40L interaction regulates T-cell tolerance, peripheral T-cell homeostasis, and T-cell-mediated inflammatory diseases.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** DKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSL SLSPGKDDDDKVTARRLNCVKHTYPSGHKCCRECPGHGMVSRCDHTRDTLCHPCETGFYNEAVNYDTCKQ CTQCNHRSGSELKQNCTPTQDTVCRCPGTQPRQDSGYKLGVCVPCPPGHFSPGNQACKPWTNCTLSGK QTRHPASDSLDAVCEDRSLLATLLWETQRPTFRPTTVQSTTVWPRTSELPSPPTLVTPPEGP

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.