

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8638: Recombinant Mouse Chordin-Like Protein 2/CHL2/CHRDL2 (C-6His)

Gene ID: Chrdl2
Gene ID: 69121
Uniprot ID: Q8VEA6

Description

Source: Human Cells.

MW:46.3kD.

Recombinant Mouse Chordin-like protein 2 is produced by our Mammalian expression system and the target gene encoding Gln24-Leu426 is expressed with a 6His tag at the C-terminus. Mouse Chordin-Like 2, also known as CHL2, is a novel chordin family member with structural homology to CHL1 which is implicated in tumor angiogenesis. The mouse CHL2 gene encodes a 426 amino acids (aa) protein with a 25 aa signal peptide. The mature chain of CHL2 protein contains two potential N-linked glycosylation sites, one putative NLS and three 63 aa cysteine-rich von Willebrand type C repeats (CRs). CHL2 gene is weakly expressed in the liver and kidney, partly expressed in the connective tissues of reproductive organs such as ligaments of the ovary and oviduct in females, and of testis, epididymis and certain male accessory sex glands in males. Recombinant mCHL2 protein interacted directly with five BMPs and one GDF thereby inhibiting, in vitro, several BMP/GDF-dependent processes including, osteogenic differentiation of C2C12 mesenchymal progenitor cells by several BMPs, ATDC5 embryonal carcinoma cells by GDF5 and BMP4-dependent lymphohematopoietic (CD34+CD31hi and CD34+CD31lo) progenitor cell development from ES cells. CHL2 may inhibits BMPs activity by blocking their interaction with their receptors, and has a negative regulator effect on the cartilage formation/regeneration from immature mesenchymal cells, by preventing or reducing the rate of matrix accumulation. Also, it may play a role during myoblast and osteoblast differentiation, and maturation.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: QARSRSGKVCLFGEKIYTPGQSWHPYLEPQGTIYCVRCTCSENGHVNCYRLRCPPLHCSQPVMEPQQCCPRCV

DPHVPSGLRVPLKSCQLNETTYQHGEIFSAQELFPARLSNQCVLCSCIEGHTYCGLMTCPEPSCPTTLPLPDSCC QTCKDRTTESSTEENLTQLQHGERHSQDPCSERRGPSTPAPTSLSSPLGFIPRHFQSVGMGSTTIKIILKEKHKK ACTHNGKTYSHGEVWHPTVLSFGPMPCILCTCIDGYQDCHRVTCPTQYPCSQPKKVAGKCCKICPEDEAEDDH SEVISTRCPKVPGQFHVYTLASPSPDSLHRFVLEHEASDQVEMYIWKLVKGIYHLVQIKRVRKQDFQKEAQNFRL

LTGTHEGYWTVFLAQTPELKVTASPDKVTKTLVDHHHHHH

Application Note

Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.