

## 32-8643: Recombinant Human Epidermal Growth Factor Receptor (V30G, Del31-297)/ErbB1/HER1 (C-6His)

**Gene :** EGFR  
**Gene ID :** 1956  
**Uniprot ID :** P00533

### Description

Source: Human Cells.

MW :39.7kD.

Recombinant Human EGFR is produced by our Mammalian expression system and the target gene encoding Leu25-Val30Gly&Asn298-Ser645 is expressed with a 6His tag at the C-terminus. The EGFR subfamily of receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43% - 44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glycoproteins with an extracellular ligand binding domain. The extracellular ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF-alpha, betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** LEEKKGNYVVTDHGSCVRACGADSYEMEEDGVRKCKKCEGPCRKVCNGIGIGEFKDSLINATNIKHFKNCTSI  
SGDLHILPVAFRGDSFTHPPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRKQHGQFSLAVV  
SLNITSLGLRSLKEISDGDVIISGNKNLCYANTINWKKLFGTSGQTKIISNRGENSCKATGQVCHALCSPEGCW  
GPEPRDCVSCRNVSRGRECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGP  
HCVKTCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPGPKIPSVDDHHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.