

32-8649: Recombinant Mouse Intercellular Adhesion Molecule 2/ICAM-2/CD102 (C-Fc)

Gene : Icam2
Gene ID : 15896
Uniprot ID : P35330

Description

Source: Human Cells.
MW :49.7kD.

Recombinant Mouse Intercellular adhesion molecule 2 is produced by our Mammalian expression system and the target gene encoding Lys23-Gln222 is expressed with a Fc tag at the C-terminus. ICAM-2 is a 55-65 kD transmembrane glycoprotein possessing 2 extracellular Ig domains, a single transmembrane domain, and a short 26-amino acid cytoplasmic domain. ICAM-2 is expressed on most leukocytes, and is strongly expressed on vascular endothelial cells. Interactions of ICAM-2 and the integrin receptors mediate cell adhesion in a wide range of lymphocyte, monocyte, natural killer cell, and granulocytewith other cells, and play important roles in many adhesion-dependent immune and inflammation responses, such as T cell aggregation, NK-cell cytotoxicity and migration, lymphocyte recirculation, etc. Serum levels of ICAM-2 correlated significantly with the inflammatory and course sequences of trichinosis in mice and had a similar relation with blood eosinophilia. So, estimation of ICAM-2 serum levels may prove useful in diagnosis of trichinosis recent infections, and in monitoring the prognosis and response to treatment.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl,pH8.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : KAFEVYIWSEKQIVEATESWKINCSTNCAAPDMGGLETPTNKIMLEEHPQGKWKQFLVSNVSKDVTFFCHFTCS GKQHSESLNIRVYQPPAQVTLKQPPRVFVGEDFTIECTVSPVQPLERLTLSELLRGRET LKNQTFGGAETVPQEA TATFNSTALKKDGLNFSCQAELDLRPHGGYIIRSISEYQILEVYEPMQDNQVDDIEGRMDEPKSCDKTHTCPPCP APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV EWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.