

32-8655: Recombinant Mouse B7-1/CD80 (C-Fc)

Gene : Cd80
Gene ID : 12519
Uniprot ID : Q00609

Description

Source: Human Cells.
MW :50.8kD.

Recombinant Mouse Activation B7-1 antigen is produced by our Mammalian expression system and the target gene encoding Val38-Lys245 is expressed with a Fc tag at the C-terminus. Cluster of Differentiation 80, also called B7-1, is a member of cell surface immunoglobulin superfamily which plays key, yet distinct roles in the activation of T cells. It is the ligand for two different proteins on the T cell surface: CD28 and CTLA-4. Studies have shown that CTLA-4 binds mostly to CD80. The structure presents two extracellular domains: a membrane distal variable-like domain (IgV) and a membrane proximal Ig constant-like domain (IgC) along with an intracellular domain. Both IgV and IgC consist of anti-parallel beta sandwiches joined by a short linker region. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKVLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLS DRGTYSVCVVQKKERGTYEVKHLALVKSIAKADFSTPNITESGNPSADTKRITCFASGGFPKPRFSWLENGRELPG INTTISQDPESELYTISSQLDFNTRNHTIKCLIKYGDAHVSEDFTWEKPPEDPPDSKVDDIEGRMDEPKSCDKTH TCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTTTPVLDSDGSSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG K

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.