

## 32-8655: Recombinant Mouse B7-1/CD80 (C-Fc)

**Gene :** Cd80  
**Gene ID :** 12519  
**Uniprot ID :** Q00609

### Description

Source: Human Cells.  
MW :50.8kD.

Recombinant Mouse Activation B7-1 antigen is produced by our Mammalian expression system and the target gene encoding Val38-Lys245 is expressed with a Fc tag at the C-terminus. Cluster of Differentiation 80, also called B7-1, is a member of cell surface immunoglobulin superfamily which plays key, yet distinct roles in the activation of T cells. It is the ligand for two different proteins on the T cell surface: CD28 and CTLA-4. Studies have shown that CTLA-4 binds mostly to CD80. The structure presents two extracellular domains: a membrane distal variable-like domain (IgV) and a membrane proximal Ig constant-like domain (IgC) along with an intracellular domain. Both IgV and IgC consist of anti-parallel beta sandwiches joined by a short linker region. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKVLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLS DRGTYSVCVVQKKERGTYEVKHLALVKSIAKADFSTPNITESGNPSADTKRITCFASGGFPKPRFSWLENGRELPG INTTISQDPESELYTISSQLDFNTRNHTIKCLIKYGDAHVSEDFTWEKPPEDPPDSKVDDIEGRMDEPKSCDKTH TCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTTTPVLDSDGSSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG K

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.