

32-8657: Recombinant Mouse IL-1 Receptor-Like 1/IL-1RL1/IL-1 R4 (C-6His)

Gene : Il1rl1
Gene ID : 17082
Uniprot ID : P14719

Description

Source: Human Cells.
MW :36.6kD.

Recombinant Mouse Interleukin-1 receptor-like 1 is produced by our Mammalian expression system and the target gene encoding Ser27-Ala337 is expressed with a 6His tag at the C-terminus. ST2, also called IL-1 R4, is an Interleukin-1 receptor family glycoprotein that plays a role in Th2 immune responses. ST2 is expressed on the surface of mast cells, activated Th2 cells, macrophages, and cardiac myocytes. This receptor is very similar to the IL-1 receptor type I and the IL-18 receptor a chain in that ST2 also has three extracellular Ig domains and an intracellular Toll domain. ST2 binds IL-33, enhances inflammatory cytokines by activating nuclear factor-kB (NF-kB) and mitogen activated protein (MAP) kinases. ST2 exists as either a membrane bound form (ST2L) or as a soluble form (sST2). ST2L acts as a transmembrane signalling receptor for IL-33 by mediating the effect of IL-33 on the inflammatory process, while sST2 can suppress IL-33 activity.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SKSSWGLENEALIVRCPQRGRSTYPVEWYSDTNESIPTQKRNRIFVSRDLKFLPARVEDSGIYACVIRSPNLN
KTGYLNVTIHKKPPSCNIPDYLMYSTVIRGSDKNFKITCPTIDLYNWTAPVQWFKNCKALQEPRFRAHRSYLFIDN
VTHDDEGDYTCQFTHAENGTNYIVTATRSFTVEEKGFMSFPVITNPPYNHTMEVEIGKPASIACSACFGKGSFHL
ADVLWQINKTVVGNFGEARIQEEEGRNESSNDMDCLTSVLRITGVTEKDLSEYDCLALNLHGMIRHTIRLR
KQPSKECPSHIAVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.