

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8709: Recombinant Mouse Nogo-66 Receptor/Reticulon 4 Receptor/NgR/RTN4R (C-6His)

Gene ID: 8tn4r **Gene ID:** 65079 **Uniprot ID:** Q99PI8

Description

Source: Human Cells. MW :46.6kD.

Recombinant Mouse NgR is produced by our Mammalian expression system and the target gene encoding Cys27-Ser447 is expressed with a 6His tag at the C-terminus. Nogo Receptor (NgR) is a glycosylphosphoinositol (GPI)-anchored protein that belongs to the Nogo recptor family. Human NgR is predominantly expressed in neurons and their axons in the central nervous systems. As a receptor for myelin-derived proteins Nogo, myelin-associated glycoprotein (MAG) and myelin oligodendrocyte glycoprotein (OMG), NgR mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult central nervous system. NgR may be proposed as a potential drug target for treatment of various neurological conditions. Additionally, NgR may play a role in regulating the function of gap junctions.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: CPGACVCYNEPKVTTSCPQQGLQAVPTGIPASSQRIFLHGNRISHVPAASFQSCRNLTILWLHSNALARIDAAAF

TGLTLLEQLDLSDNAQLHVVDPTTFHGLGHLHTLHLDRCGLRELGPGLFRGLAALQYLYLQDNNLQALPDNTF RDLGNLTHLFLHGNRIPSVPEHAFRGLHSLDRLLLHQNHVARVHPHAFRDLGRLMTLYLFANNLSMLPAEVLMP LRSLQYLRLNDNPWVCDCRARPLWAWLQKFRGSSSEVPCNLPQRLADRDLKRLAASDLEGCAVASGPFRPIQ TSQLTDEELLSLPKCCQPDAADKASVLEPGRPASAGNALKGRVPPGDTPPGNGSGPRHINDSPFGTLPSSAEPP

LTALRPGGSEPPGLPTTGPRRRPGCSRKNRTRSHCRLGQAGSGASGTGDAEGSVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ \tilde{A} \parallel \hat{A} μ g (1 IEU/ \tilde{A} \parallel \hat{A} μ g) as determined by LAL test.