

32-8716: Recombinant Mouse Endothelial Cell-Specific Molecule /Endocan/ESM-1 (C-6His)

Gene : Esm1
Gene ID : 71690
Uniprot ID : Q9QYY7

Description

Source: Human Cells.
MW :19kD.

Recombinant Mouse ESM-1 is produced by our Mammalian expression system and the target gene encoding Trp20-Arg184 is expressed with a 6His tag at the C-terminus. Endothelial cell-specific molecule-1 (ESM-1)—so-called endocan—is a novel endothelium derived soluble dermatan sulfate proteoglycan (PG) that is constitutively expressed by endothelial cells in lungs and kidneys and can be detected in human blood. It is encoded by the ESM1 gene. The expression of this gene is regulated by several cytokines and growth factors, such as vascular endothelial growth factor. Inflammatory cytokines, such as interleukin (IL)-1 beta and tumor necrosis factor (TNF)- α , stimulate the upregulation of endocan mRNA and the secretion of endocan from endothelial cells. The binding of circulating endocan to leukocyte ligand for ICAM-1—Lymphocyte Function-associated Antigen-1 (LFA-1) and to leukocyte ligand for VCAM-1—Very Late Antigen-4 (VLA-4) is important in leukocyte adhesion and interaction with activated endothelium. Endocan is a key player in the regulation of major processes such as cell adhesion in inflammatory disorders and tumor progression.

Product Info

Amount : 6His)(Discontinued) / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : WSAKYAVDCPEHCDKTECRSSLRCKRTLDDCGCCQVCAAGPGETCYRTVSGMDGVKCGPLKCHFYEED DFGDEFGICKDCPYGTFGMECKETCNCQSGICDRVTGRCLDFPFQYAAAKSPRTSASHTERDSASGDGNAV REEIGEGNAARPSVMKWLNPRVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.