

32-8766: Recombinant Cynomolgus TIGIT/VSIG9/VSTM3 (C-6His)

Gene : EGM_10359

Uniprot ID : G7NXM4

Description

Source: Human Cells.

MW :14.2kD.

Recombinant Cynomolgus monkey TIGIT is produced by our Mammalian expression system and the target gene encoding Met89-Pro209 is expressed with a 6His tag at the C-terminus. T cell immunoreceptor with Ig and ITIM domains (TIGIT), also called VSIG9 and Vstm3, is a member of the CD28 family within the Ig superfamily of proteins. TIGIT contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM), and is expressed on regulatory, memory, activated T cells and NK cells. TIGIT binds to CD155(PVR) that appear on dendritic cells (DC), macrophages and endothelium with high affinity, and CD112(PVRL2) with lower affinity, but not CD113 (PVRL3). TIGIT-Fc fusion protein could interact with PVR on DC and enhance the secretion of IL-10, but inhibit the macrophage activation. Mice lacking TIGIT show increased T cell responses and susceptibility to autoimmune challenges, while knockdown of TIGIT with siRNA in human memory T cells did not affect T cell responses.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : MMTGTIETTGNISAKKGGSVILQCHLSSTMAQVTQVNWEQHDHSLLAIRNAELGWHIYPAFKDRVAPGPGGLGL
TLQSLTMNDTGEYFCTYHTYPDGTGRIFLEVLESSVAEHSARFQIPHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.