

32-8794: Recombinant Human IL-1 Receptor Type 2/IL-1R-2 (C-6His)

Gene : IL1R2
Gene ID : 7850
Uniprot ID : P27930

Description

Source: Human cells.

MW :38.6kD.

Recombinant Human Interleukin-1 receptor type 2/IL-1R-2 is produced by our Mammalian expression system and the target gene encoding Phe14-Glu343 is expressed fused with a 6His tag at the C-terminus. Interleukin-1 receptor type 2 (IL1R2) belongs to the interleukin-1 receptor family. Two distinct types of IL1 receptors which are able to bind IL1 specifically have been identified, designated as IL1RI (IL1RA) and IL1RII (IL1RB). IL1 receptor type II is a 68 kDa transmembrane protein found on B lymphocytes, neutrophils, monocytes, large granular leukocytes and endothelial cells. IL1R2 is non-signaling receptor for IL1A, IL1B and IL1RN, reduces IL1B activities. IL1R2 serves as a decoy receptor by competitive binding to IL1B and preventing its binding to IL1R1. IL1R2 modulates cellular response through non-signaling association with IL1RAP after binding to IL1B. IL1R2 (membrane and secreted forms) preferentially binds IL1B and poorly IL1A and IL1RN. The secreted IL1R2 recruits secreted IL1RAP with high affinity; this complex formation may be the dominant mechanism for neutralization of IL1B by secreted/soluble receptors.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : FTLQPAAHTGAARSCFRGRHYKREFLEGE PVALRCPQVPYWLWASVSPRINLTWHKND SARTVPGEETR
MWAQDGALWLLPALQEDSGTYVCTTRNASYCDKMSIELRVFENTDAFLPFISYPQILTLSTSGVLVCPDLSEFTR
DKTQDKIQWYKDSLDDKNEKFLSVRGTTLLVHDVALEDAGYYRCVLTFAHEGQQYNITRSIELRIKKKEETI
PVIISPLKTISASLGSRILTIPCKVFLGTGTPLTTLWWTANDTHIESAYPGGRVTEGPRQEYSENNENYIEVPLIFD
PVTREDLHMDFKCVVHNTLSFQTLRTTVKEHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.