

32-8801: Recombinant Human Ephrin A Receptor 2/EphA2 (C-Fc)

Gene : EPHA2
Gene ID : 1969
Uniprot ID : P29317

Description

Source: Human Cells.
MW :83kD.

Recombinant Human Ephrin A Receptor 2 is produced by our Mammalian expression system and the target gene encoding Ala24-Asn534 is expressed with a Fc tag at the C-terminus. Ephrin type-A receptor 2/EphA2 is a member of the Eph receptor tyrosine kinase family which binds Ephrins A1, 2, 3, 4, and 5. A and B class Eph proteins have a common structural organization. Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. EphA2 becomes autophosphorylated following ligand binding and then interacts with SH2 domain-containing PI3-kinase to activate MAPK pathways. Reverse signaling is also propagated through the Ephrin ligand. Transcription of EphA2 is dependent on the expression of E-Cadherin, and can be induced by p53 family transcription factors. EphA2 is upregulated in breast, prostate, and colon cancer vascular endothelium. Its ligand, EphrinA1, is expressed by the local tumor cells. In some cases, EphA2 and EphrinA1 are expressed on the same blood vessels. EphA2 signaling cooperates with VEGF receptor signaling in promoting endothelial cell migration.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AQQKEVLLDFAAAGGELGWLTHPYGKGWDLMQNIMNDMPIYMYVCNVMSGDQDNWLRNWWVYRGEAE RIFIELKFTVRDCNSFPGGASSCKETFNLYAESDLDTGTFNQKRLFTKIDTIAPDEITVSSDFEARHVKLNVEERS VGPLTRKGFYLAHQDIGACVALLSVRVYKCKPELLOGLAHFPETIAGSDAPSLATVAGTCVDHAVVPPGGEEPR MHCAVDGEWLVPIGQCLCQAGYEKVEDACQACSPGFFKFEASESPCLECPEHTLPSPEGATSCCEEGFFRAP QDPASMPCTRPPSAPHYLTAVGMGAKVELRWTPPQDSGGREDIVYSVTCEQCWPESGECGPCEASVRYSEPP HGLTRTSVTVSDLEPHMNYTFTVEARNGVSLVTSRFSRTASVSINQTEPPKVRLEGRSTTSLSVSWSIPPPQQS RVWKYEVTYRKKGDSNSYNVRRTEGFSVTLDDLAPDTTYLVQVQALTQEGQGAGSKVHEFQTLSPGSGNDI EGRMDPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMHEA LHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.