

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8803: Recombinant Mouse Thrombopoietin/THPO/TPO (N-6His)

Gene ID: Thpo
Gene ID: 21832
Uniprot ID: P40226

## **Description**

Source: Human Cells. MW :36.4kD.

Recombinant Mouse Thrombopoietin is produced by our expression system and the target gene encoding Ser22ÂThr356 is expressed with a 6His tag at the N-terminus. Thrombopoietin (TPO) is a glycoprotein hormone which belongs to the EPO/TPO family. It produced by the liver and kidney which regulates the production of platelets. Mature mouse Tpo shares 71% and 81% aa sequence homology with human and rat Tpo, respectively. It is an 80-85 kDa protein that consists of an N-terminal domain with homology to Erythropoietin (Epo) and a C-terminal domain that contains multiple N-linked and O-linked glycosylation sites. TPO stimulates the production and differentiation of megakaryocytes, the bone marrow cells that bud off large numbers of platelets. Lineage-specific cytokine affects the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.

## **Product Info**

**Amount:** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: HHHHHHSPVAPACDPRLLNKLLRDSHLLHSRLSQCPDVDPLSIPVLLPAVDFSLGEWKTQTEQSKAQDILGAVS

LLLEGVMAARGQLEPSCLSSLLGQLSGQVRLLLGALQGLLGTQLPLQGRTTAHKDPNALFLSLQQLLRGKVRFL LLVEGPTLCVRRTLPTTAVPSSTSQLLTLNKFPNRTSGLLETNFSVTARTAGPGLLSRLQGFRVKITPGQLNQTSR SPVOISGYLNRTHGPVNGTHGLFAGTSLOTLEASDISPGAFNKGSLAFNLQGGLPPSPSLAPDGHTPFPPSPALP

TTHGSPPQLHPLFPDPSTTMPNSTAPHPVTMYPHPRNLSQET

## **Application Note**

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.