

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8872: Recombinant Human Apoptosis Regulator Bcl-2 (C-6His)(Discontinued)

Gene ID: 596 **Uniprot ID:** P10415

Description

Source: E.coli. MW:24.1kD.

Recombinant Human Apoptosis Regulator Bcl-2 is produced by our E.coli expression system and the target gene encoding Met1-Asp211 is expressed with a 6His tag at the C-terminus. Bcl-2 is a member of a family of proteins that regulates outer mitochondrial membrane permeability. Bcl-2 is an antiapoptotic member that prevents release of cytochrome c from the mitochondria intermembrane space into the cytosol. Bcl-2 is present on the outer mitochondrial membrane and is also found on other membranes in some cell types. BCL-2 is localized to the outer membrane of mitochondria, where it plays an important role in promoting cellular survival and inhibiting the actions of pro-apoptotic proteins. The pro-apoptotic proteins in the BCL-2 family, including Bax and Bak, normally act on the mitochondrial membrane to promote permeabilization and release of cytochrome C and ROS, that are important signals in the apotosis cascade. These pro-apoptotic proteins are in turn activated by BH3-only proteins, and are inhibited by the function of BCL-2 and its relative BCL-XI.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content :Supplied as a 0.2 μm filtered solution of 20mM HEPES, 150mM NaCl, 10%Glycerol, pH8.0.Storage condition :Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.

Amino Acid: MAHAGRTGYDNREIVMKYIHYKLSQRGYEWDAGDVGAAPPGAAPAPGIFSSQPGHTPHPAASRDPVARTSPL

QTPAAPGAAAGPALSPVPPVVHLTLRQAGDDFSRRYRRDFAEMSSQLHLTPFTARGRFATVVEELFRDGVNW GRIVAFFEFGGVMCVESVNREMSPLVDNIALWMTEYLNRHLHTWIQDNGGWDAFVELYGPSMRPLFDHHHH

НН

Application Note

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.