

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8886: Recombinant Human GDF-5/BMP-14

Gene ID: 8200 **Uniprot ID:** P43026

Description

Source: E. coli. MW :13.7kD.

Recombinant Human Growth/Differentiation Factor 5 is produced by our E.coli expression system and the target gene encoding Ala382-Arg501 is expressed. Growth Differentiation Factor 5(GDF-5, BMP-14) is a member of the BMP family of TGF beta superfamily proteins. Human GDF-5, -6, and -7 are a defined subgroup of the BMP family. GDF-5 is synthesized as a homodimeric precursor protein consisting of a 354 amino acid (aa) Nterminal proregion and a 120 aa C-terminal mature peptide. Mature human GDF-5 shares 99% aa sequence identity with both mature mouse and rat GDF-5. GDF-5 signaling is mediated by formation of a heterodimeric complex consisting of a type 1 (BMPR-IB) and a type II (BMPR-IIor Activin RII) serine/threonine kinase receptor which results in the phosphorylation and activation of cytosolic Smad proteins (Smad1, 5, and 8). GDF-5 is involved in multiple developmental processes including limb generation, cartilage development, joint formation, bone morphogenesis, cell survival, and neuritogenesis. Inhibition of GDF-5 expression or alteration of its signaling can facilitate the development of osteoarthritis.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 4mM HCl.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MAPLATROGKRPSKNLKARCSRKALHVNFKDMGWDDWIIAPLEYEAFHCEGLCEFPLRSHLEPTNHAVIQTLM

NSMDPESTPPTCCVPTRLSPISILFIDSANNVVYKQYEDMVVESCGCR

Application Note

Endotoxin : Less than $0.1 \text{ ng/}\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.