

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8888: Recombinant Human Olfactomedin-4/OLFM4 (C-10His)

Gene ID: 0LFM4
Gene ID: 10562
Uniprot ID: Q6UX06

## **Description**

Source: Human Cells. MW :56.9kD.

Recombinant Human Olfactomedin-4 is produced by our Mammalian expression system and the target gene encoding Asp21-Gln510 is expressed with a 10His tag at the C-terminus. Olfactomedin-4/OLFM4 is a secreted protein which contains one olfactomedin-like domain. OLFM4 is expressed during myeloid lineage development, it strongly expressed in the prostate, small intestine, colon and moderately expressed in the bone marrow and stomach. OLFM4 is an antiapoptotic factor that promotes tumor growth. It expressed at high levels in stomach cancer and colon cancer tissues. it promotes proliferation of pancreatic cancer cells by favoring the transition from the S to G2/M phase. In myeloid leukemic cell lines, OLFM4 inhibits cell growth and induces cell differentiation and apoptosis. Through interaction with cell surface lectins and cadherin, OLFM4 facilitates cell adhesion. It may play a role in the inhibition of EIF4EBP1 phosphorylation/deactivation. Induction of OLFM4 in cancer cells was reported to have a novel antiapoptotic action via binding to the potent apoptosis inducer GRIM-19.

## **Product Info**

**Amount:** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: DLGDVGPPIPSPGFSSFPGVDSSSSFSSSRSGSSSSRSLGSGGSVSQLFSNFTGSVDDRGTCQCSVSLPDTTF

PVDRVERLEFTAHVLSQKFEKELSKVREYVQLISVYEKKLLNLTVRIDIMEKDTISYTELDFELIKVEVKEMEKLVIQ LKESFGGSSEIVDQLEVEIRNMTLLVEKLETLDKNNVLAIRREIVALKTKLKECEASKDQNTPVVHPPPTPGSCGH GGVVNISKPSVVQLNWRGFSYLYGAWGRDYSPQHPNKGLYWVAPLNTDGRLLEYYRLYNTLDDLLLYINAREL RITYGQGSGTAVYNNNMYVNMYNTGNIARVNLTTNTIAVTQTLPNAAYNNRFSYANVAWQDIDFAVDENGLW VIYSTEASTGNMVISKLNDTTLQVLNTWYTKQYKPSASNAFMVCGVLYATRTMNTRTEEIFYYYDTNTGKEGKL

DIVMHKMQEKVQSINYNPFDQKLYVYNDGYLLNYDLSVLQKPQGGGGSHHHHHHHHHH

## **Application Note**

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \sqcap \hat{A} \mu g$  (1 IEU/ $\tilde{A} \sqcap \hat{A} \mu g$ ) as determined by LAL test.