

32-8909: Recombinant Cynomolgus Fibroblast Growth Factor 21/FGF-21 (C-6His)

Description

Source: Human Cells.

MW :20.2kD.

Recombinant Cynomolgus Fibroblast Growth Factor 21 (Accession No.-XM_005589811.2) is produced by our Mammalian expression system and the target gene encoding His29-Ser209 is expressed with a 6His tag at the C-terminus. Fibroblast Growth Factor 21 (FGF21) is a growth factor that belongs to the FGF family. FGF family proteins play a central role during prenatal development and postnatal growth and regeneration of many tissues, by promoting cellular proliferation and differentiation. FGF21 is a potent activator of glucose uptake on adipocytes, protects animal from diet-induced obesity when overexpression in transgenic mice, and lower blood glucose and triglyceride levels when therapeutically administered to diabetic rodents. FGF21 is produced by hepatocytes in response to free fatty acid stimulation of a PPAR α /RXR dimeric complex. This situation occurs clinically during starvation, or following the ingestion of a highly-fat/low-carbohydrate diet. Upon FGF21 secretion, white adipose tissue is induced to release free fatty acids from triglyceride stores. Once free fatty acids reach hepatocytes, they are oxidized and reduced to acetyl-CoA. The acetyl-CoA is recombined into 4-carbon ketone bodies, release, and transported to peripheral tissue for TCA processing and energy generation.

Product Info

Amount : 10 μ g / 50 μ g

Content : Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : HPIPDSSPLLQFGGQVRQRYLYTDDAQQTEAHLEIREDDGTVGGAHQSPESLLQLKALKPGVIQILGVKTSRFLCQKPDGALYGSLHFDPEACSFRELLLENGYNVYQSEAHGLPLHLPGNKSPHRDPASQGPAPFLPLPGLPPAPPEP
PGILAPQPPDVGSSDPLSMVGPSQARSPSYASHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.