

## 32-8923: Recombinant Human Interleukin-4 Receptor Subunit Alpha/IL-4 Ra(C-Fc)

**Gene :** IL4R  
**Gene ID :** 3566  
**Uniprot ID :** P24394

### Description

Source: Human cells.  
MW :50.2kD.

Recombinant Human Interleukin-4 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Met26-Gln231 is expressed with a Fc tag at the C-terminus. Interleukin 4 Receptor alpha (IL4-Ra) is a widely expressed 140 kDa transmembrane glycoprotein in the class I cytokine receptor family. Mature human IL4-Ra consists of a 207 amino acid (aa) extracellular domain (ECD) that contains a cytokine binding region and one fibronectin type III domain, a 24 aa transmembrane segment, and a 569 aa cytoplasmic domain that contains one Box 1 motif and one ITIM motif. IL4-Ra plays an important role in Th2-biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. Soluble forms of IL4-Ra, generated by alternate splicing or proteolysis, retain ligand binding properties and inhibit IL-4 bioactivity. IL4-Ra is a component of two distinct receptor complexes and shows species selectivity between human and mouse. It can associate with the common gamma chain ( gamma c) to form the IL-4 responsive type I receptor in which gamma c increases the affinity for IL-4 and enables signaling. It can alternatively associate with IL13-Ra1 to form the type II receptor which is responsive to both IL-4 and IL-13. The use of shared receptor components contributes to the overlapping biological effects of IL-4 and IL-13 as well as other cytokines that utilize gamma c.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MKVLQEPTECVSDYMSISTCEWKMNGPTNCSTELRLLYQLVFLLEAHTCIPENNGGAGCVCHLLMDDVVSADN  
YTLDLWAGQQLLWKGSEFKPSEHVKPRAPGNLTVHTNVSDTLTLLTWSNPYPPDNYLYNHLYAVNIWSENDPA  
DFRIYNVTYLEPSLRIAASTLKSIGSYRARVRAWAQCYNNTTWESEWSPSTKWHNSYREPFEQIEGRMDPKSCDKT  
HTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN  
STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGF  
YPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVDFSCVMHEALHNHYTQKSLSLSP  
GK

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.