

## 32-8927: Recombinant Human Macrophage Colony-stimulating Factor 1 Receptor/M-CSF R/CSF1R/CD115(C-Fc)

**Gene :** CSF1R  
**Gene ID :** 1436  
**Uniprot ID :** P07333

### Description

Source: Human cells.

MW :81.9kD.

Recombinant Human Macrophage Colony-stimulating Factor 1 Receptor is produced by our Mammalian expression system and the target gene encoding Ile20-Pro517 is expressed with a Fc tag at the C-terminus. Macrophage colony-stimulating factor 1 receptor (CSF1R) is a member of the type III subfamily of receptor tyrosine kinases that also includes receptors for SCF and PDGF. These receptors each contain five immunoglobulin-like domains in their extracellular domain (ECD) and a split kinase domain in their intracellular region. CSF1R is expressed primarily on cells of the monocyte/macrophage lineage, dendritic cells, stem cells and in the developing placenta. CSF1 and its receptor (CSF1R, product of c-fms proto-oncogene) were initially implicated as essential for normal monocyte development as well as for trophoblastic implantation. It plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone and tooth development. It is required for normal male and female fertility, and for normal development of milk ducts and acinar structures in the mammary gland during pregnancy. Aberrant expression of CSF1 or CSF1R may play a role in inflammatory diseases, such as rheumatoid arthritis, glomerulonephritis, atherosclerosis, and allograft rejection.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** IPVIEPSVPELVVKPGATVTLRCVGNCSVEWDGPPSPHWTLYSYGSSSILSTNNATFQNTGTYRCTEPGDPLGG SAAIHLYVKDPPARPWNVLAQEVEVFEDQDALLPCLLTDPVLEAGVSLVRVGRPLMRHTNYSFSPWHGFTIHR AKFIQSQDYQCSALMGGRKVMISIRLKVQKVIPGPPALTLVPAELVRIRGEAAQIVCSASSVDVNFDFLQHNNTKLAIPQQSDFHNNRYQKVLTLNLDQVDFQHAGNYSCVASNVQGHSTSMFFRVVESAYLNLSSSEQNLIQEVTVGEGNLKVMVEAYPGLQGFNWTYLGPFSDHQPEPKLANATTKD TYRHTFTLSLPRLKPSEAGRYSFLARNPG GWRALTFELTLRYPPEVSVIWFINGSGTLLCAASGYQPNTWLQCSGHTDRCDQAQVLQVWDDPYEVLSQ EPFHKVTVQSLLTVETLEHNQTYECRAHNSVSGSWAFIPI SAGAHTHPDEF LFTPDIEGRMDPKSCDKHTC PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSV MHEALHNHYTQKLSLSLSPGK

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.