

32-8931: Recombinant Mouse Inducible T-cell Costimulator/ICOS(C-6His)

Gene : Icos
Gene ID : 54167
Uniprot ID : Q9WVS0

Description

Source: Human cells.
MW :14.7kD.

Recombinant Mouse Inducible T-cell Costimulator is produced by our Mammalian expression system and the target gene encoding Glu21-Leu142 is expressed with a 6His tag at the C-terminus. Inducible Costimulator(ICOS) is a member of the growing CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA4 and PD1. ICOS shares approximately 39% amino acid similarity with CD 28 and CTLA4. Mouse and human ICOS share approximately 72% amino acid identity. ICOS is expressed on most CD45RO+ cells. ICOS expression is up-regulated within approximately 24-48 hours of activation on Th primed cells. B7-H2, a member of the B7 family of costimulatory ligands, has been identified as the ICOS ligand. The B7-H2/ ICOS interaction appears to play roles in T cell dependent B cell activation and Th differentiation. In addition, ICOS is more potent in the induction of IL-10 production, acytokine important for suppressive function of T regulatory cells.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : EINGSADHRMFSFHNGGVQISCKYPETVQQLKMRLFREREVLCELTCTKSGNAVSIKPNMLCLYHLSNNSVSF
FLNNPDSSQGSYYFCSLSIFDPPPFQERNLSGGYLHIYESQLCCQLKLHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.