

## 32-8932: Recombinant Human Cytotoxic T-lymphocyte Protein 4(C-Flag)

**Gene :** CTLA4  
**Gene ID :** 1493  
**Uniprot ID :** P16410

### Description

Source: Human cells.  
MW :14.5kD.

Recombinant Human Cytotoxic T-lymphocyte Protein 4 is produced by our Mammalian expression system and the target gene encoding Lys36-Asp161 is expressed with a Flag tag at the C-terminus. Cytotoxic Tlymphocyte 4(CTLA-4,CD152), is a type I transmembrane T cell inhibitory molecule that is a member of the Ig superfamily. Human or mouse CTLA4 cDNA encodes 223 amino acids (aa) including a 35 aa signal sequence, a 126 aa extracellular domain (ECD) with one Ig-like V-type domain, a 21 aa transmembrane (TM) sequence, and a 41 aa cytoplasmic sequence. It is widely expressed with highest levels in lymphoid tissues. CD28 and CTLA-4, together with their ligands, B7-1 and B7-2, constitute one of the dominant costimulatory pathways that regulate T and B cell responses. CD28 and CTLA-4 are structurally homologous molecules that are members of the immunoglobulin (Ig) gene superfamily. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T Cells and may play an important role in their functions. Tcell activation through the Tcell receptor and CD28 leads to increased expression of CTLA4.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** KAMHVAQPAVVCLASSRGIA SFVCEYASPGKATEVRVTVL RQADSQVTEVCAATYMMGNELTFLDDSICTGTSS  
GNQVNLTIQGLRAMDTGLYICKVELMYPPPYL GIGNGTQIYVIDPEPCPDSDDYKDDDDK

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.