

32-8941: Recombinant Human Hepatocyte Growth Factor Receptor/HGF R/cMet (C-6His)

Gene : MET
Gene ID : 4233
Uniprot ID : P08581

Description

Source: Human Cells.
MW :102.5kD.

Recombinant Human Hepatocyte Growth Factor Receptor is produced by our Mammalian expression system and the target gene encoding Glu25-Thr932 is expressed with a 6His tag at the C-terminus. Hepatocyte growth factor receptor (HGF R) is a glycosylated receptor tyrosine kinase that plays a central role in epithelial morphogenesis and cancer development. HGF R is synthesized as a single chain precursor which undergoes cotranslational proteolytic cleavage. Mature HGF R is a disulfide-linked dimer composed of a 50 kDa extracellular α chain and a 145 kDa transmembrane β chain. Proteolysis and alternate splicing generate additional forms of human HGF R which either lack of the kinase domain, consist of secreted extracellular domains, or are deficient in proteolytic separation of the α and β chains. The sema domain, which is formed by both α and β chains of HGF R, mediates both ligand binding and receptor dimerization. HGF stimulation induces HGF R downregulation via internalization and proteasomedependent degradation. Paracrine induction of epithelial cell scattering and branching tubulogenesis results from the stimulation of HGF R on undifferentiated epithelium by HGF released from neighboring mesenchymal cells.

Product Info

Amount : 10 μ g / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.
Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : ECKEALAKSEMNVNMKYQLPNFTAETPIQNVILHEHHIFLGATNYIYVLNEEDLQKVAEYKTPVLEHPDCFCPCQDCSSKANLSSGGVWKDNINMALVVDTYDDQLISCGSVNRGTCQRHVFPNHHTADIQSEVHCIFSPQIEEPSQCPCCVVSALGAKVLSSVKDRFINFFVGNTINSSYFPDHLHSISVRRKTKDGFMLTDQSYIDVLPFRDSDYPIKYVHAFESNNFIYFLTVQRETLDAQTFHTRIIRFCSINSGLHSEMPLECILTEKRKKRSTKKEVFNILQAAVSKPGAQLARQIGASLNDDILFGVFAQSKPDSAEPMDRSAMCAFPKIYVNDFFNKIVNKNVRCLOHFYGPNEHCFNRTLNRSSGCEARRDEYRTEFTALQRVDLFGQFSEVLLTSISTFIKGDILTIANLGTSEGRFMQVVSRSGPSTPHVNFLLDSDHPVSPEVIVEHTLNQNGYTLVITGKKITKIPLNGLGCRHFQSCSQCLSAPPFVQCGWCHDKCVRSEECLSGTWTQQICLPAIYKVPNSAPLEGGTRLTICGWDFGFRNKNKFDLKKTRVLLGNESCTLTLESTMTNLKCTVGPAMNKHFNMSIIISNGHGTQYSTFSYVDPVITSISPKYGPAGGTLTGTNYLNSGNSRHISIGGKTCTLKSVSNSILECYTPAQTISTEFAVKKIDLANRETSIFSREDPIVYIEIHPTKSFISGGSTITGVGKNLNSVSVPRMVINVHEAGRNFTVACQHRNSSEIICCTPSLQQLNLQLPLKTKAFFMLDGILSKYFDLIYVHNPVFKPFKPEKPMISMGNENVLEIKGNDIDPEAVKGEVLKVGKSCENIHLHSEAVLCTVPNDLLKLNSELNIEWKQAISSVTLGKVIVQPDQNFTHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.