

32-8979: Recombinant Human LILRA5/CD85f (C-6His)

Gene :	LILRA5
Gene ID :	353514
Uniprot ID :	A6NI73

Description

Source: Human Cells.

MW :26.1kD.

Recombinant Human Leukocyte Immunoglobulin-like Receptor Subfamily A Member 5 is produced by our Mammalian expression system and the target gene encoding Gly42-Arg268 is expressed with a 6His tag at the C-terminus. Leukocyte Immunoglobulin-like Receptor Subfamily A Member 5(LILRA5) is a member of the leukocyte immunoglobulin-like receptors (LILR), comprise a family of activating and inhibitory type immunoreceptors. LILRA5 consists of a 227 amino acid (aa) extracellular domain (ECD), a 21 aa transmembrane segment, and a 10 aa cytoplasmic tail. The ECD contains two Ig-like domains and the transmembrane segment contains a positively charged aspartic acid residue which may mediate its association with the signaling molecule, FcR common gamma chain. LILRA5 is expressed by monocytes, macrophages, and neutrophils. Cross-linking of LILRA5 on monocytes induces the expression of pro-inflammatory cytokines (IL-1beta, IL-6, TNF-alpha) as well as the anti-inflammatory IL-10. It can be detected in tissues of the hematopoietic system, including bone marrow, spleen, lymph node and peripheral leukocytes. Crosslink of ILT-11 on the surface of monocytes has been shown to induce calcium flux and secretion of several proinflammatory cytokines, which suggests the roles of this protein in triggering innate immune responses.

Product Info

Amount :	10 μg / 50 μg
Content :	Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	GNLSKATLWAEPGSVISRGNSVTIRCQGTLEAQEYRLVKEGSPEPWDTQNPLEPKNKARFSIPSMTEHHAGRYR CYYYSPAGWSEPSDPLELVVTGFYNKPTLSALPSPVVTSGENVTLQCGSRLRFDRFILTEEGDHKLSWTLDSQLT PSGQFQALFPVGPVTPSHRWMLRCYGSRRHILQVWSEPSDLLEIPVSGAADNLSPSQNKSDSGTASHLQDYAV ENLIRHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ \tilde{A} $\hat{A}\mu g$ (1 IEU/ \tilde{A} $\hat{A}\mu g$) as determined by LAL test.