

32-8983: Recombinant Human Cytotoxic T-lymphocyte Protein 4 (C-Fc-Avi)

Gene : CTLA4
Gene ID : 1493
Uniprot ID : P16410

Description

Source: Human Cells.
MW :41.5kD.

Recombinant Human Cytotoxic T-lymphocyte Protein 4 is produced by our Mammalian expression system and the target gene encoding Lys36-Asp161 is expressed with a Fc & Avi tag at the C-terminus. Cytotoxic T-lymphocyte protein 4, is a single-pass type I membrane protein. It is widely expressed with highest levels in lymphoid tissues. CD28 and CTLA-4, together with their ligands, B7-1 and B7-2, constitute one of the dominant costimulatory pathways that regulate T and B cell responses. CD28 and CTLA-4 are structurally homologous molecules that are members of the immunoglobulin (Ig) gene superfamily. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T Cells and may play an important role in their functions. Tcell activation through the Tcell receptor and CD28 leads to increased expression of CTLA4.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : KAMHVAQPAVVCLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATYMMGNETLFLDDSICTGTSS
GNQVNLTIQGLRAMDTGLYICKVELMYPYPYLGIGNGTQIYVIDPEPCPDSQEPKSSDKTHTSPSPAPPELLG
GSSVFLFPPKPKDTLMISRTPVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLH
QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESN
GQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCVMHEALHNHYTQKSLSLSPGKGLNDIFEAQK
IEWHE

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.