

32-8997: Recombinant Human Tyrosine-protein Kinase Receptor TYRO3 (C-6His)

Gene : TYRO3
Gene ID : 7301
Uniprot ID : Q06418

Description

Source: Human Cells.
MW :42.4kD.

Recombinant Human Tyrosine-protein Kinase Receptor TYRO3 is produced by our Mammalian expression system and the target gene encoding Ala41-Ser428 is expressed with a 6His tag at the C-terminus. Axl (Ufo, Ark), Dtk (Sky, Tyro3, Rse, Brt) and Mer (human and mouse homologues of chicken cEyk) constitute a new receptor tyrosine kinase subfamily. The extracellular domain of these proteins contain two Ig-like motifs and two fibronectin type III motifs. This characteristic topology is also found in neural cell adhesion molecules and in receptor tyrosine phosphatases. All three receptors bind the vitamin K-dependent protein growth-arrest specific gene 6 (Gas6) which is structurally related to the anticoagulation factor protein S. The binding affinities for Gas6 is in the order of Axl > Dtk > Mer. Gas6 binding induces tyrosine phosphorylation and downstream signaling pathways that can lead to cell proliferation, migration, or the prevention of apoptosis. Dtk is widely expressed during embryonic development. In adults, Dtk is predominantly expressed in neurons in restricted regions of the brain.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AGLKLMGAPVKLTVSQGPVKLNCSVEGMEEPDIQWVKDGA VVQNL DQLYIPVSEQHWIGFLSLKSVERSDA
GRYWCQVEDGGETEISQPVWLTVEGV PFFTVEPKDLAVPPNAPFQLSCEAVGPPPEVTIVVWRGTTKIGGPAP
SPSVLNVTGVTQSTMFSC EAHNLKGLASSRTATVHLQALPAAPFNITVTKLSSSNASVAWMPGADGRALLQSC
TVQVTQAPGGWEVLAVVVPVPPFTCLLRDLVPATNYS LRVCANALGSPYADWVPFQTKGLAPASAPQNLHA
IRTDSGLILEWEEVIPEAPLEGPLGPYKLSWVQDNGTQDEL TVEGTRANLTGWDPQKDLIVRVCVSNVAVGCGP
WSQPLVSSSHDRAGQQGPPHSRTSHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.