

## 32-9023: Recombinant Human Angiopoietin-like Protein 3/ANGPTL3 (C-6His)

**Gene :** ANGPTL3

**Gene ID :** 27329

**Uniprot ID :** Q9Y5C1

### Description

Source: Human Cells.

MW :24.6kD.

Recombinant Human Angiopoietin-related Protein 3 is produced by our expression system and the target gene encoding Ser17-Pro220 is expressed. Angiopoietin-like 3 (ANGPTL3) is a secreted glycoprotein that is structurally related to the angiopoietins. Mature human ANGPTL3 contains an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain. ANGPTL3 is expressed in the liver from early in development through adulthood. Full length ANGPTL3 circulates in the plasma as do the proteolytically separated N- and C-terminal segments containing the coiled coil domain and fibrinogen-like domains, respectively. ANGPTL3 directly inhibits lipoprotein lipase (LPL) and endothelial lipase (EL), enzymes responsible for hydrolyzing circulating triglycerides and HDL phospholipids. ANGPTL3 promotes an increase in circulating triglyceride levels without altering VLDL or HDL secretion or uptake. ANGPTL3 expression in vivo is up-regulated by LXR agonists and down-regulated by insulin, leptin, and agonists of TR beta or PPAR beta.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** SRIDQDNSSFDSLSPKSRFAMLDDVKILANGLLQLGHGLKDFVHKTGQINDIFQKLNIFDQSFYDLSLQTSEIKEEKELRRTTYKLQVKNEEVKNMSLELNSKLESLLKILLQKVKYLEEQLTNLIQNQPETPEHPEVTSKTFVEKQDNSIKDLLQTVEDQYKQLNQHSQIKEIENQLRRTSIQEPTEISLSSKPHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.