

## 32-9027: Recombinant Human C-type lectin domain family 4 member K/CD207 (N-6His)(Discontinued)

**Gene :** CD207  
**Gene ID :** 50489  
**Uniprot ID :** Q9UJ71

### Description

Source: Human Cells.

MW :31.5kD.

Recombinant Human C-type lectin domain family 4 member K is produced by our expression system and the target gene encoding Tyr64-Pro328 is expressed Langerin (CD207) is a type II transmembrane glycoprotein which is member K of the C-type lectin domain family. Langerin is used as a marker for Langerhans cells (LCs) which represent the immature dendritic cells in the epidermis. Langerin is necessary and sufficient for Birbeck granule formation. Human langerin sequence contains a 43 aa cytoplasmic domain, a 21 aa transmembrane domain and a 264 aa extracellular domain (ECD) that contains a coiled-coil domain and a single C-type lectin domain. Human langerin shares 68%, 62%, 71% aa identity with mouse, rat and bovine langerin ECD, respectively. Trimerization greatly increases the lectin binding affinity. Langerin internalizes endogenous proteins such as type I procollagen. Internalization by LC is thought to lead to suppression of self reactions. Langerin also mediates endocytosis of non-peptide antigens containing mannose, N-acetyl glucosamine and fucose that are expressed by mycobacteria and fungus.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** GSGSHHHHHHIEGRYPRFMGTISDVKTNVQLLGRVDNISTLDSEIKKNSDGMEEAGVQIQMVNESLGIVRSQ  
FLKLTTSVEKANAQIQLTRSWEEVSTLNAQIPELKSLEKASALNTKIRALQGSLENMSKLLKRQNDILQVVSQG  
WKYFKGNFYFSLIPKTWYSAEQFCVSRNSHLTSVTSESEQEFLYKTAGGLIYWIGLTKAGMEGDWSWVDDTP  
FNKVQSARFWIPGEPNAGNNEHCGNIKAPSLQAWNDAPCDKTFLEFICKRYPVVPSEP

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.