

32-12029: Human Epstein-Barr Virus Induced Gene 3 Subunit (AF)

Gene : EBI3
Gene ID : 10148
Uniprot ID : Q14213
Alternative Name : Epstein-Barr virus-induced gene 3 protein, IL27B

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 23.4 kDa (210 aa)

Epstein-Barr virus induced gene-3 (EBI3) is a secreted glycoprotein belonging to the hematopoietin receptor family related to the p40 subunit of interleukin 12 (IL-12). EBI3 expression is induced in B-lymphocytes in response to Epstein-Barr virus infection. EBI3 forms heterodimers with p28 to form interleukin 27 (IL-27), and with p35 to form interleukin 35 (IL-35). Both IL-27 and IL-35 have anti-inflammatory and regulatory activity.

Product Info

Amount : 20 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 90\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) + 0.5% mannitol
Sterile 10 mM HCl at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MRKGPPAALT LPRVQCRASR YPIAVDCSWT LPPAPNSTSP VSFATYRLG MAARGHSWPC LQQTPTSTSC
TITDVQLFSM APYVLNVTAV HPWGSSSSSFV PFITEHIKP DPPEGVRLSP LAERQLQVQW EPPGSWPFPE
IFSLKYWIRY KRQGAARFHR VGPIEATSF I LRAVRPRARY YVQVAAQDLT DYGELSDWSL PATATMSLGK

Application Note

Endotoxin: Less than 0.5 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.

